arXiv:1211.6555v1 [math.AC] 28 Nov 2012

Vertex covers and sensor networks

MAURIZIO IMBESI
Faculty of Sciences, Department of Mathematics,
University of Messina
Viale Ferdinando Stagno d'Alcontres, 31 − I 98166 Messina, Italy
e-mail: imbesim@unime.it

MONICA LA BARBIERA
Faculty of Sciences, Department of Mathematics,
University of Messina
Viale Ferdinando Stagno d'Alcontres, 31 − I 98166 Messina, Italy
e-mail: monicalb@unime.it

Abstract

We consider algebraic developments of graph theory through suitable applications in real connection problems. We investigate ideals of vertex covers for the edge ideals associated to a significative class of connected graphs. It is shown that algebraic procedures linked to minimal vertex covering of such graphs are good instruments concerned about how sensor networks can be encoded and their properties measured. Moreover algebraic properties of such ideals are studied. Using the notion of linear quotients, standard algebraic invariants are computed and these ideals are stated to be Cohen-Macaulay.

AMS 2010 Subject Classification: 05C25, 05E40, 13C15.

Key words and phrases: Ideals of vertex covers. Sensor network. Linear quotients and standard invariants.

Introduction

In this paper we are interested in studying standard properties of monomial ideals arising from a class of graphs in order to explain how algebraic models, built through graph theory, are useful for setting suitable solutions of several problems concerning network areas (infrastructure nets, circuits electrical workers, urban and territorial analysis, wireless sensor networks, etc...).
In the last years, Commutative and Combinatorial Algebra have paid particular attention to large classes of monomial ideals derived from graphs in order to describe algebraic properties of them. Such ideals turn out to be

very interesting from a combinatorial point of view too. Classical topics include graph algebraic theory, in particular the edge ideals, namely monomial ideals generated in degree two which descend from the edges of a graph. Moreover, using the notion of minimal covering for the nodes of a graph, recent results show that it is possible to attach to graphs the so-called ideals of vertex covers; an intensive study of them carries to solve interesting and unthinkable problems related to these fields. More precisely, the ideals of vertex covers of a wide class of simple graphs are examined and useful properties are studied. Moreover, we explain how algebraic methods arising from graph theory are suitable solutions for several problems concerning the *wireless sensor networks*.

Wireless sensor networks (WSNs) are an important technology for large-scale monitoring, providing sensor measurements at high temporal and spatial resolution. A sensor network is composed of a large number of sensor nodes, which are densely deployed either inside the phenomenon or very close to it. These sensor nodes, which consist of sensing, data processing, and communicating components, leverage the idea of sensor networks based on collaborative effort of a large number of nodes. Sensor nodes are fitted with an on-board processor. Instead of sending the raw data to the nodes responsible for the fusion, sensor nodes use their processing abilities to locally carry out simple computations and transmit only the required and partially processed data. The above described features ensure a wide range of monitoring applications for sensor networks. Some of the application areas are environmental, military, security and commercial ([1]).

We consider the problem of the placement of as few as possible leader nodes which collect the information received by the sensors and send them to a base station, so that each sensor is near to at least one of them. This problem is related to the energy saving of the sensors. The suitable model to represent this situation is a simple graph $\mathcal{G}$ and the problem is linked to the study of the minimal vertex covers of $\mathcal{G}$. We introduce a class of simple graphs and some algebraic aspects linked to the minimal vertex covers are analyzed. We show that the algebraic procedures introduced are good instruments concerned how this problem related to wireless sensor networks can be encoded and solved.

Let $\mathcal{G}$ be a graph on n vertices $v_1, \ldots, v_n$. An algebraic object attached to $\mathcal{G}$ is the edge ideal $I(\mathcal{G})$, a monomial ideal of the polynomial ring in n variables $R = K[X_1, \ldots, X_n]$, K a field.

When $\mathcal{G}$ is a simple graph, $I(\mathcal{G})$ is generated by squarefree monomials of degree two in R, $I(\mathcal{G}) = \big(\{X_iX_j \mid \{v_i, v_j\} \text{ is an edge of } \mathcal{G}\}\big)$.

The *ideal of vertex covers* of $I(\mathcal{G})$, denoted by $I_c(\mathcal{G})$ is the ideal of R generated by all the monomials $X_{i_1} \cdots X_{i_k}$ such that $(X_{i_1}, \ldots, X_{i_k})$ is an associated prime ideal of $I(\mathcal{G})$ ([9]). Under special situations, $I(\mathcal{G})$ reflects properties of $I_c(\mathcal{G})$ and the study of $I(\mathcal{G})$ states some properties of $I_c(\mathcal{G})$. In particular, the Cohen-Macaulay property of the ideals $I_c(\mathcal{G})$ was studied by G. Lyubeznik in [8] using graph theoretical methods; the central notion is that of chordal graph, that is the right notion to express some homological properties of edge ideals and the Cohen-Macaulay property of the ideal of vertex covers.

We examine a wide class of squarefree edge ideals associated to connected graphs $\mathcal{H}$ consisting of the union of a complete graph K_m and star graphs with centers the vertices of K_m. The ideals of vertex covers for the edge ideals associated to the these graphs were described in [6]. The generators of these ideals give all the minimal vertex covers and their cardinality. If $\mathcal{H}$ is considered to be the model of a WSN, then the vertex covers with minimum cardinality give the position of the leaders in the sensor network that $\mathcal{H}$ represents.

Moreover we study some algebraic aspects of the ideals $I_c(\mathcal{H})$. We prove that $I_c(\mathcal{H})$ has linear quotients. In general, if a monomial ideal of R generated in one degree has linear quotients, then it has a linear resolution ([2], Lemma 4.1). In our case, $I_c(\mathcal{H})$ has linear quotients, but in general $I_c(\mathcal{H})$ has not a linear resolution because it is not generated in the same degree. Using some properties of the graph $\mathcal{H}$ and of its edge ideal we can obtain conditions such that $I_c(\mathcal{H})$ has a linear resolution and some algebraic invariants are computed. Starting from theoretic properties of the graph $\mathcal{H}$ and of its edge ideal we can obtain information about the Cohen-Macaulay property of $I(\mathcal{H})$ and of its ideal of vertex covers.

The work is subdivided as follows. In section 1 we describe the developments of graph theory to networks concerning of problems of minimal vertex covers. The class of graphs $\mathcal{H}$ and the ideals of vertex covers $I_c(\mathcal{H})$ are analyzed as a geometric model of a WSN using computational and algebraic methods. In section 2 we examine algebraic properties of $I_c(\mathcal{H})$. We prove that these ideals have linear quotients and we investigate conditions such that they have a linear resolution. Moreover using the technique of studying the linear quotients of such ideals, as previously employed in [7], standard algebraic invariants are calculated.

1 Developments of graph theory to networks

The aim of this section is to explain how algebraic and geometric models, built through graph theory, are useful for setting suitable solutions of several problems concerning the *wireless sensor networks.*

A wireless sensor network (WSN) consists of spatially distributed autonomous sensors (called sensor nodes) to monitor physical or environmental conditions (such as temperature, sound, vibration, pressure, humidity,...) and to pass their data through the network to a main location. The sensors collect data of interest and send them to special gateway nodes, which collect the information received by the sensors and send them to a base station. The sensor networks can improve the quality of information: for example, can guarantee a high fidelity, can provide information in real time and can reduce the cost of transmitting the same information.

Current research indicates highly developments for this type of wireless networks which are already in use in many fields. Useful applications of sensor networks are environmental, military, and commercial. Area monitoring is a common application of WSNs that are deployed over a region where some phenomenons are to be monitored: the urban traffic or natural phenomena such as glaciers, volcanoes and earthquakes. Some environmental applications of sensor networks include tracking the movements of small animals; detecting foreign chemical agents in the air and the water, irrigation; precision agriculture; environmental monitoring in marine; forest fire detection; meteorological or geophysical research. Some of the commercial applications are monitoring product quality; interactive museums; monitoring disaster area; transportation; vehicle tracking and detection ([1]).

Consider a sensor network in which each sensor must send information it has collected to the nearest gateway node. The problem is the placement of as few as possible gateway nodes so that each sensor is near to at least one of them. This problem is related to the energy saving of the sensors. The formalization of the model to represent the sensor network is given by a graph $\mathcal{G} = (V(\mathcal{G}), E(\mathcal{G}))$, $V(\mathcal{G}) = \{v_1, \ldots, v_n\}$ is the set of nodes and $E(\mathcal{G}) = \{\{v_i, v_j\} | v_i \neq v_j, \quad v_i, v_j \in V(\mathcal{G})\}$ is the set of edges (if two nodes v_1 and v_2 are adjacent then $\{v_1, v_2\} \in E(\mathcal{G})$).

Our problem consists of finding a set of nodes leaders (gateway nodes) to be responsible to send messages to nodes belonging to the higher level. Such leaders must have the feature of being near to all nodes. This means that we want to find a subset of nodes $C \subset V(\mathcal{G})$ such that each node in $V(\mathcal{G}) \setminus C$ is joined to a node in C. This is a problem of minimum vertex cover. The optimization problem of minimum vertex cover is looking for a covering of vertices of a graph that has minimum cardinality. The corresponding decision problem of vertex cover is, given a graph $\mathcal{G}$ and an integer k, if there is a vertex covering of $\mathcal{G}$ of minimal cardinality k or of cardinality at most k.

The basic network topologies are represented by complete graphs, meshes, star graphs, trees. A single network may consist of several interconnected subnets of different topologies. We consider as a network topology a class of simple graphs union of a complete graph and star graphs, and some algebraic

aspects linked to this problem of minimal vertex covers are analyzed.
Let's introduce some preliminary notions.
Let $\mathcal{G}$ be a graph. A *cycle* of $\mathcal{G}$ of length q is a subgraph C_q of $\mathcal{G}$ such that $E(C_q) = \{\{v_{i_1}, v_{i_2}\}, \{v_{i_2}, v_{i_3}\}, \ldots, \{v_{i_{q-1}}, v_{i_q}\}, \{v_{i_q}, v_{i_1}\}\}$, where $\{v_{i_1}, \ldots, v_{i_q}\} \in V(\mathcal{G})$ and $v_{i_j} \neq v_{i_k}$ if $i_j \neq i_k$. A *chord* of a cycle C_q is an edge $\{v_{i_j}, v_{i_k}\}$ of $\mathcal{G}$ such that v_{i_j} and v_{i_k} are vertices of C_q with $\{v_{i_j}, v_{i_k}\} \notin E(C_q)$. A graph $\mathcal{G}$ is said *chordal* or *triangulated* if every cycle C_q in $\mathcal{G}$, $q > 3$, has a chord.
A graph $\mathcal{G}$ on n vertices $v_1, \ldots, v_n$ is *complete* if there exists an edge for all pairs $\{v_i, v_j\}$ of vertices of $\mathcal{G}$. It is denoted by K_n.
A *star graph* on vertex set $\big\{\{v_i\}, \{v_1, \ldots, v_{i-1}, v_{i+1}, v_n\}\big\}$ with center v_i, denoted by $\mathrm{star}_i(n)$, $i = 1, \ldots, n$, is a complete bipartite graph of the form $K_{1,n-1}$. The *complementary graph* of a graph $\mathcal{G}$ on $\{v_1, \ldots, v_n\}$ is the graph $\overline{\mathcal{G}}$ on $\{v_1, \ldots, v_n\}$ whose edge set $E(\overline{\mathcal{G}})$ consists of those 2-element subset $\{v_i, v_j\}$ for which $\{v_i, v_j\} \notin E(\mathcal{G})$.

If $V(\mathcal{G}) = \{v_1, \ldots, v_n\}$ and $R = K[X_1, \ldots, X_n]$ is the polynomial ring over a field K such that each variable X_i corresponds to the vertex v_i, the *edge ideal* $I(\mathcal{G})$ associated to $\mathcal{G}$ is the ideal $\big(\{X_iX_j \mid \{v_i, v_j\} \in E(\mathcal{G})\}\big) \subset R$.

Definition 1.1. Let $\mathcal{G}$ be a graph with vertex set $V(\mathcal{G}) = \{v_1, \ldots, v_n\}$. A subset C of $V(\mathcal{G})$ is said a *minimal vertex cover* for $\mathcal{G}$ if:
(1) every edge of $\mathcal{G}$ is incident with one vertex in C;
(2) there is no proper subset of C with this property.

If C satisfies condition (1) only, then C is called a *vertex cover* of $\mathcal{G}$ and C is said to cover all the edges of $\mathcal{G}$. The smallest number of vertices in any minimal vertex cover of $\mathcal{G}$ is said *vertex covering number*. We denote it by $\alpha_0(\mathcal{G})$.
We consider some algebraic aspects linked to the minimal vertex covers of a graph $\mathcal{G}$ with set of edges $E(\mathcal{G})$. More precisely, starting by the notion of minimal vertex cover, recent results show that is possible to attach to graphs the so-called *ideals of vertex covers* ([9]). There exists a one to one correspondence between the minimal vertex covers of $\mathcal{G}$ and the minimal prime ideals of $I(\mathcal{G})$. In fact, $\wp$ is a minimal prime ideal of $I(\mathcal{G})$ if and only if $\wp = (C)$, for some minimal vertex cover C of $\mathcal{G}$ ([9], Prop. 6.1.16). Hence $\mathrm{ht}(I(\mathcal{G})) = \alpha_0(\mathcal{G})$. Thus the primary decomposition of the edge ideal of $\mathcal{G}$ is given by $I(\mathcal{G}) = (C_1) \cap \cdots \cap (C_p)$, where $C_1, \ldots, C_p$ are the minimal vertex covers of $\mathcal{G}$.

Definition 1.2. The *ideal of vertex covers* of $I(\mathcal{G})$, denoted by $I_c(\mathcal{G})$, is the ideal of R generated by all monomials $X_{i_1} \cdots X_{i_k}$ such that $(X_{i_1}, \ldots, X_{i_k})$ is an associated prime ideal of $I(\mathcal{G})$.

Hence $I_c(\mathcal{G}) = (\{X_{i_1}\cdots X_{i_k} \,|\, \{v_{i_1},\dots,v_{i_k}\}$ is a vertex cover of $\mathcal{G}\})$ and the minimal generators of $I_c(\mathcal{G})$ correspond to the minimal vertex covers.

Now we study this topic for a wide class of connected graphs and we show how it is possible to apply the results to the problem related to wireless sensor networks. More precisely, $\mathcal{H}$ is the connected graph on n vertices $v_1,\dots,v_n$ consisting of the union of a complete graph K_m and star graphs in the vertices of K_m: $\mathcal{H} = K_m \cup \operatorname{star}_j(k)$, where K_m is the complete graph on m vertices $v_1,\dots,v_m$, $m < n$, and $\operatorname{star}_j(k)$ is the star graph on k vertices with center v_j, for some $j = 1,\dots,m$, $k \leq n-m$. We observe that $V(\operatorname{star}_j(k)) \cap V(K_m) = \{v_j\}$ for $j = 1,\dots,m$. We denote $|E(\operatorname{star}_j(k))| = i_j$, for $j = 1,\dots,m$. The edge ideal $I(\mathcal{H}) \subset R = K[X_1,\dots,X_n]$ is stated in [6]: $I(\mathcal{H}) = (X_1X_{m+1}, X_1X_{m+2},\dots,X_1X_{m+i_1}, X_1X_2, X_2X_{m+i_1+1}, X_2X_{m+i_1+2},\dots,$ $X_2X_{m+i_1+i_2}, X_2X_3, X_1X_3, X_3X_{m+i_1+i_2+1}, X_3X_{m+i_1+i_2+2},\dots, X_3X_{m+i_1+i_2+i_3},$ $X_3X_4,\ X_2X_4, X_1X_4,\dots, X_mX_{m+i_1+i_2+\cdots+i_{m-1}+1}, X_mX_{m+i_1+i_2+\cdots+i_{m-1}+2},\dots,$ $X_mX_{m+i_1+i_2+\cdots+i_{m-1}+i_m})$.

Remark 1.1. $I(\mathcal{H})$ has a linear resolution ([6], Prop 1.1). R. Fröberg proved that the edge ideal of a simple graph has a linear resolution if and only if its complementary graph is chordal ([3]; [4], Th. 9.2.3). Hence $\overline{\mathcal{H}}$ is chordal.

In [6] it is given the structure of the ideal of vertex cover of $I(\mathcal{H})$. In order to describe $I_c(\mathcal{H})$ we rename the vertices of the complete graph K_m $v_{\alpha_1},\dots,v_{\alpha_m}$, $1 \leqslant \alpha_1 < \alpha_2 < \dots < \alpha_m = n$, and $\alpha_1 = 1$ if there is not the star graph with center v_{α_1}.

The following proposition gives explicitly the generators of the ideals of vertex covers for the edge ideals $I(\mathcal{H})$ when at least a vertex of K_m has degree $m-1$.

Proposition 1.1. *([6], Prop. 2.1)*
Let $\mathcal{H} = K_m \cup star_j(k)$ with n vertices $v_1,\dots,v_n$ and $K_m, m < n$, with vertices $v_{\alpha_1},\dots,v_{\alpha_m}$, $1 \leqslant \alpha_1 < \alpha_2 < \dots < \alpha_m = n$. Let $star_j(k)$ be of the type $star_{\alpha_1}(\alpha_1)$ on vertices $v_1,\dots,v_{\alpha_1}$, or $star_{\alpha_i}(\alpha_i - \alpha_{i-1})$, for some $i = 2,\dots,m$. Then $I_c(\mathcal{H})$ has m monomial squarefree generators and it is:
$(X_1\cdots X_{\alpha_1-1}X_{\alpha_2}\cdots X_{\alpha_m}, X_{\alpha_1}X_{\alpha_3}\cdots X_{\alpha_m},\dots,X_{\alpha_1}\cdots X_{\alpha_{m-1}})$, *if* $i = 1$,
or
$(X_{\alpha_2}\cdots X_{\alpha_m},\dots,X_{\alpha_1}\cdots X_{\alpha_{i-1}}X_{\alpha_{i-1}+1}\cdots X_{\alpha_{i+1}-1}X_{\alpha_{i+1}}\cdots X_{\alpha_m},\dots,X_{\alpha_1}\cdots X_{\alpha_{m-1}})$,
otherwise.

More in general, it is given the structure of the ideal of vertex covers of $I(\mathcal{H})$ when all the vertices of K_m have degree at least m.

Theorem 1.1. *([6], Th. 2.1)*
Let $\mathcal{H} = K_m \cup star_j(k)$ with m star graphs with centers each of the vertices of K_m. The ideal of vertex covers of $I(\mathcal{H})$ has $m+1$ monomial squarefree generators and $I_c(\mathcal{H})$ is
$(X_{\alpha_1}X_{\alpha_2}\cdots X_{\alpha_m}, X_1\cdots X_{\alpha_1-1}X_{\alpha_2}\cdots X_{\alpha_m}, X_{\alpha_1}X_{\alpha_1+1}\cdots X_{\alpha_2-1}X_{\alpha_3}\cdots X_{\alpha_m}, \dots, X_{\alpha_1}\cdots X_{\alpha_{m-2}}X_{\alpha_{m-2}+1}\cdots X_{\alpha_{m-1}-1}X_{\alpha_m}, X_{\alpha_1}\cdots X_{\alpha_{m-1}}X_{\alpha_{m-1}+1}\cdots X_{\alpha_m-1})$.

The monomial ideal $I_c(\mathcal{H})$ is generated in degree $\geq m$. Hence using the previous results we give the answer to the *decision problem of vertex cover*: given the graph $\mathcal{H}$ there is a vertex covering of $\mathcal{H}$ of minimum cardinality m, where m is the number of vertices of the complete graph K_m. Moreover the generators of $I_c(\mathcal{H})$ give the structure of each vertex covering of $\mathcal{H}$.

Example 1.1. Let $\mathcal{H}$ be the simple graph with $V(\mathcal{H}) = \{v_1, \dots, v_{11}\}$

$$\mathcal{H} = K_4 \cup \text{star}_2(2) \cup \text{star}_6(4) \cup \text{star}_8(2) \cup \text{star}_{11}(3).$$

This graph is the model of a wireless sensor network.



$I_c(\mathcal{H})$ is generated by $\{X_1X_6X_8X_{11}, X_2X_6X_8X_{11}, X_2X_6X_7X_{11}, X_2X_6X_8X_9X_{10}, X_2X_3X_4X_5X_8X_{11}\}$. The minimal vertex covers of $\mathcal{H}$ are:
$C_1 = \{v_1, v_6, v_8, v_{11}\}$, $C_2 = \{v_2, v_6, v_8, v_{11}\}$, $C_3 = \{v_2, v_6, v_7, v_{11}\}$,
$C_4 = \{v_2, v_6, v_8, v_9, v_{10}\}$, $C_5 = \{v_2, v_3, v_4, v_5, v_8, v_{11}\}$.
Choose $C_1 = \{v_1, v_6, v_8, v_{11}\}$ among the three vertex covers with minimum cardinality. C_1 gives the position of the leaders in the sensor network that $\mathcal{H}$ represents:



2 Algebraic properties of $I_c(\mathcal{H})$

The class of monomial ideals $I_c(\mathcal{H})$ is a suitable algebraic model to solve problems of minimal vertex covers. Moreover $I_c(\mathcal{H})$ is an interesting class of monomial ideals for its good algebraic properties. In this section we study some of them.

Let $R = K[X_1, \ldots, X_n]$ be the polynomial ring in n variables over a field K with $\deg X_i = 1$, for all $i = 1, \ldots, n$. For a monomial ideal $I \subset R$ we denote by $G(I)$ its unique minimal set of monomial generators.

Definition 2.1. A monomial ideal $I \subset R$ has *linear quotients* if there is an ordering $u_1, \ldots, u_t$ of monomials belonging to $G(I)$ such that the colon ideal $(u_1, \ldots, u_{j-1}) : (u_j)$ is generated by a subset of $\{X_1, \ldots, X_n\}$, for $2 \leqslant j \leqslant t$.

For a monomial ideal I of R having linear quotients with respect to the ordering $u_1, \ldots, u_t$ of the monomials of $G(I)$, let $q_j(I)$ denote the number of the variables which is required to generate the ideal $(u_1, \ldots, u_{j-1}) : (u_j)$.

Set $q(I) = \max_{2 \leqslant j \leqslant t} q_j(I)$. The integer $q(I)$ is independent on the choice of the ordering of the generators that gives linear quotients ([5]).

Proposition 2.1. *$I_c(\mathcal{H})$ has linear quotients.*

Proof. 1) If at least a vertex of K_m has degree $m-1$ (Proposition 1.1), then $I_c(\mathcal{H})$ has m monomial squarefree generators. We have the following cases:

a) $\mathcal{H}$ is formed by the union of a complete graph $K_m, m < n$, with vertices $v_{\alpha_1}, \ldots, v_{\alpha_m}$, $1 \leqslant \alpha_1 < \alpha_2 < \ldots < \alpha_m = n$, and of a star graph $\mathrm{star}_{\alpha_1}(\alpha_1)$ on vertices $v_1, \ldots, v_{\alpha_1}$. Then we set the following order for the monomial generators of $I_c(\mathcal{H})$: $f_1 = X_{\alpha_1} \cdots X_{\alpha_{m-1}}$, $f_2 = X_{\alpha_1} \cdots X_{\alpha_{m-2}} X_{\alpha_m}$, $\ldots$, $f_{m-1} = X_{\alpha_1} X_{\alpha_3} \cdots X_{\alpha_m}$, $f_m = X_1 \cdots X_{\alpha_1 - 1} X_{\alpha_2} \cdots X_{\alpha_m}$.

It results: $(f_1) : (f_2) = (X_{\alpha_{m-1}})$, $(f_1, f_2) : (f_3) = (X_{\alpha_{m-2}})$, $\ldots$, and so on up to $(f_1, \ldots, f_{m-1}) : (f_m) = (X_{\alpha_1})$.

b) $\mathcal{H}$ is formed by the union of a complete graph $K_m, m < n$, with vertices $v_{\alpha_1}, \ldots, v_{\alpha_m}$, $1 \leqslant \alpha_1 < \alpha_2 < \ldots < \alpha_m = n$, and of a star graph $\mathrm{star}_{\alpha_i}(\alpha_i - \alpha_{i-1})$, for some $i = 2, \ldots, m$. Then we set the following order for the monomial generators of $I_c(\mathcal{H})$: $f_1 = X_{\alpha_1} \cdots X_{\alpha_{m-1}}$, $f_2 = X_{\alpha_1} \cdots X_{\alpha_{m-2}} X_{\alpha_m}$, $\ldots$, $f_{m-1} = X_{\alpha_1} X_{\alpha_2} \cdots X_{\alpha_m}$, $f_m = X_{\alpha_2} \cdots X_{\alpha_m}$. Hence: $(f_1) : (f_2) = (X_{\alpha_{m-1}})$, $(f_1, f_2) : (f_3) = (X_{\alpha_{m-2}})$, $\ldots$, and so on up to $(f_1, \ldots, f_{m-1}) : (f_m) = (X_{\alpha_1})$.

2) If all the vertices of K_m have degree at least m (Theorem 1.1), then we set the following order for the monomial generators of $I_c(\mathcal{H})$: $f_1 = X_{\alpha_1} X_{\alpha_2} \cdots X_{\alpha_m}$, $f_2 = X_{\alpha_1} \cdots X_{\alpha_{m-1}} X_{\alpha_{m-1}+1} \cdots X_{\alpha_m - 1}$, $f_3 = X_{\alpha_1} \cdots X_{\alpha_{m-2}} X_{\alpha_{m-2}+1} \cdots X_{\alpha_{m-1}-1} X_{\alpha_m}$, $\ldots$, $f_m = X_{\alpha_1} X_{\alpha_1+1} \cdots X_{\alpha_2 - 1} X_{\alpha_3} \cdots X_{\alpha_m}$, $f_{m+1} = X_1 \cdots X_{\alpha_1 - 1} X_{\alpha_2} \cdots X_{\alpha_m}$. It results: $(f_1) : (f_2) = (X_{\alpha_m})$, $(f_1, f_2) : (f_3) = (X_{\alpha_{m-1}})$, $(f_1, f_2, f_3) : (f_4) = (X_{\alpha_{m-2}})$,

$\ldots$, and so on up to $(f_1, \ldots, f_m) : (f_{m+1}) = (X_{\alpha_1})$.
Hence $I_c(\mathcal{H})$ has linear quotients. ☐

Remark 2.1. If a monomial ideal I of R generated in one degree has linear quotients, then I has a linear resolution ([2], Lemma 4.1). $I_c(\mathcal{H})$ has linear quotients, but in general it has not a linear resolution because it is not generated in the same degree.

Using some properties of the graph $\mathcal{H}$ and of its edge ideal $I(\mathcal{H})$ we can obtain the conditions for which $I_c(\mathcal{H})$ has a linear resolution.

Theorem 2.1. *(([6], Th. 1.1)*
Let $R = K[X_1, \ldots, X_n]$ and $I(\mathcal{H}) \subset R$, $\mathcal{H} = K_m \cup$ $star_j(k)$. Let $|E(star_j(k))| = i_j$, for $j = 1, \ldots, m$. Then:

1) $dim_R(R/I(\mathcal{H})) = n - m$.

2) $pd_R(R/I(\mathcal{H})) = m + max_{1\leqslant j\leqslant m}\, i_j - 1$.

3) $depth_R(R/I(\mathcal{H})) = n - m - max_{1\leqslant j\leqslant m}\, i_j + 1$.

4) $reg_R(R/I(\mathcal{H})) = 1$.

Proposition 2.2. *Let $I(\mathcal{H}) \subset R = K[X_1, \ldots, X_n]$. $I(\mathcal{H})$ is Cohen Macaulay if and only if $k = 2$.*

Proof. Let $I(\mathcal{H})$ be the edge ideal of the graph $\mathcal{H}$ as described in section 1. By Theorem 2.1 one has $\dim_R(R/I(\mathcal{H})) = n-m$ and $\operatorname{depth}_R(R/I(\mathcal{H})) = n-m-\max_{1\leqslant j\leqslant m} i_j + 1$. $I(\mathcal{H})$ is Cohen Macaulay if and only if $\dim_R(R/I(\mathcal{H})) = \operatorname{depth}_R(R/I(\mathcal{H}))$. Hence the equality holds if and only if $\max_{1\leqslant j\leqslant m} i_j = 1$. This is equivalent to say that $|E(\text{star}_j(k))| = 1$ for all $j = 1, \ldots, m$, i.e. $\text{star}_j(k)$ has $k = 2$ vertices for all $j = 1, \ldots, m$. ☐

Proposition 2.3. *Let $\mathcal{H} = K_m \cup$ $star_j(k)$. Then $I_c(\mathcal{H})$ has linear resolution if and only if $k = 2$.*

Proof. $I(\mathcal{H})$ has linear resolution ([6], Prop. 1.1), that is equivalent to say that the complementary graph of $\mathcal{H}$ is chordal ([4], Th. 9.2.3). By Proposition 2.2, $k = 2$ if and only if $I(\mathcal{H})$ is Cohen-Macaulay. Hence by [9] (Prop. 6.7.15) this is equivalent to say that $I_c(\mathcal{H})$ has linear resolution. ☐

In general for a graph $\mathcal{G}$ one has $I_c(\mathcal{G}) = \bigcap\limits_{\substack{X_iX_j \in I(\mathcal{G})\\ i\neq j}} (X_i, X_j)$.

Hence $I_c(\mathcal{G})$ is an unmixed ideal of height two ([9]). But in general it is not Cohen-Macaulay. The Cohen-Macaulay property of the ideal $I_c(\mathcal{G})$ was studied in [8]. The notion of chordal graph is used to express this algebraic property and to study an interesting duality between $I(\mathcal{G})$ and $I_c(\mathcal{G})$ ([3]). For the class of graphs $\mathcal{H}$ starting by some properties of the graph $\mathcal{H}$ and of its edge ideal, we state information about the Cohen-Macaulay property of $I_c(\mathcal{H})$.

Proposition 2.4. *$I_c(\mathcal{H})$ is Cohen-Macaulay.*

Proof. $I(\mathcal{H})$ has linear resolution ([6], Prop. 1.1). Hence the complementary graph $\overline{\mathcal{H}}$ of $\mathcal{H}$ is chordal ([4], Th. 9.2.3). This is equivalent to say that $I_c(\mathcal{H})$ is Cohen-Macaulay ([8]). $\square$

The study of the linear quotients is useful in order to investigate algebraic invariants of $R/I_c(\mathcal{H})$: the dimension, $\dim_R(R/I_c(\mathcal{H}))$, the depth, $\operatorname{depth}(R/I_c(\mathcal{H}))$, the projective dimension, $\operatorname{pd}_R(R/I_c(\mathcal{H}))$.

Lemma 2.1. *Let $R = K[X_1, \dots, X_n]$ and $I_c(\mathcal{H}) \subset R$. Then $q(I_c(\mathcal{H})) = 1$.*

Proof. By the computation of the linear quotients (Proposition 2.1), the number of the variables which is required to generate the ideal $(f_1, \dots, f_{h-1}) : (f_h)$ is 1, for all $h = 1, \dots, t$, $t \leq m+1$. $\square$

Theorem 2.2. *Let $R = K[X_1, \dots, X_n]$ and $I_c(\mathcal{H}) \subset R$. Then:*

1) $pd_R(R/I_c(\mathcal{H})) = 2$

2) $depth_R(R/I_c(\mathcal{H})) = n-2$

3) $dim_R(R/I_c(\mathcal{H})) = n-2$

4) $reg_R(R/I_c(\mathcal{H})) = max\{deg(v_{\alpha_i})\} - 1$, $i = 1, \dots, m$.

Proof. 1) The length of the minimal free resolution of $R/I_c(\mathcal{H})$ over R is equal to $q(I_c(\mathcal{H})) + 1$ ([5], Corollary 1.6). Then $\operatorname{pd}_R(R/I_c(\mathcal{H})) = 2$.
2) As a consequence of 1), by Auslander-Buchsbaum formula, one has $\operatorname{depth}_R(R/I_c(\mathcal{H})) = n - \operatorname{pd}_R(R/I_c(\mathcal{H})) = n-2$.
3) By Proposition 2.4 $I_c(\mathcal{H})$ is Cohen-Macaulay, hence $\dim_R(R/I_c(\mathcal{H})) = \operatorname{depth}_R(R/I_c(\mathcal{H})) = n-2$.
4) By [2] (Lemma 4.1) the Castelnuovo-Mumford regularity is $\operatorname{reg}_R(R/I_c(\mathcal{H})) = \max\{\deg(f) | \ f$ minimal generator of $I_c(\mathcal{H})\} - 1$. By the structure of generators of $I_c(\mathcal{H})$ it follows that $\max\{\deg(f) | \ f$ minimal generator of $I_c(\mathcal{H})\} = m - 1 - \max\{\deg(v_{\alpha_i})\} - (m-1) = \max\{\deg(v_{\alpha_i})\}$, $i = 1, \dots, m$. Hence $\operatorname{reg}_R(R/I_c(\mathcal{H})) = \max\{\deg(v_{\alpha_i})\} - 1$, $i = 1, \dots, m$. $\square$

Example 2.1. Let $\mathcal{H} = K_3 \cup \text{star}_1(2) \cup \text{star}_2(2) \cup \text{star}_3(2)$ on vertices $\{v_1, \ldots, v_6\}$.
$R = K[X_1, \ldots, X_6]$, $I(\mathcal{H}) = (X_1X_2, X_2X_6, X_2X_4, X_4X_6, X_3X_4, X_5X_6)$.
$I_c(\mathcal{H}) = (X_2X_4X_6, X_2X_4X_5, X_2X_3X_6, X_1X_4X_6)$ is generated in the same degree 3. $I_c(\mathcal{H})$ has linear quotients, hence linear resolution:

$$0 \to R^3(-4) \to R^4(-3) \to R \to R/I_c(\mathcal{H}) \to 0$$

One has:

1) $\text{pd}_R(R/I_c(\mathcal{H})) = 2$.

2) $\dim_R(R/I_c(\mathcal{H})) = \text{depth}_R(R/I_c(\mathcal{H})) = 4$.

3) $\text{reg}_R(R/I_c(\mathcal{H})) = 2$

References

[1] Akyildiz I.F., Su W., Sankarasubramaniam Y., Cayirci E. - Wireless sensor networks: a survey. *Computer Networks* **38** (2002), 393-422.

[2] Conca, A., and Herzog, J. - Castelnuovo-Mumford regularity of product of ideals. *Collect. Math.* **54** (2003), 137–152.

[3] Fröberg, R. - On Stanley-Reisner rings. *Topics in Algebra* (1990), 57–70.

[4] Herzog, J., and Hibi, T. - Monomial Ideals. *Springer, Graduate Texts in Mathematics* **260** (2011).

[5] Herzog, J., and Takayama, Y. - Resolutions by mapping cones. *Homology, Homotopy Appl.* **4**, 2 (2002), 277–294.

[6] Imbesi, M., and La Barbiera M. - Monomial ideals of graphs with loops. *Anal. Ştiinţ. Univ. Al. I Cuza din Iaşi, Sect. Mat.*, In press.

[7] La Barbiera M. - On a class of monomial ideals generated by s-sequences. *Math. Reports*, **12(62)**, 3 (2010), 201–216.

[8] Lyubeznik, G. - The minimal non-Cohen-Macaulay monomial ideals. *J. Pure and Applied Algebra*, **51** (1988), 261–266.

[9] Villarreal, R.H. - Monomial Algebras. *M. Dekker Inc., New York, Pure and Appl. Math.* **238** (2001).